

# DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

## Warnings

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- asthma
- taken a diuretic

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

### Other information

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see WARNINGS)
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- repeat dose every **6-8 hours**, if needed
- do not use more than **4 times a day**
- measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

#### Dosing Chart

Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1½ tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2½ tsp
72-95 lb	11 yr	3 tsp

- each teaspoon contains: sodium 3 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)

- see bottom of box for lot number and expiration date
- **each teaspoon contains:** sodium 10 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

***Inactive ingredients***

artificial flavors, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

acetic acid, artificial flavor, butylated hydroxytoluene, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C blue no.1, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

artificial and natural flavors, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C blue no. 1, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucrose, xanthan gum